



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

59. ASCOBOLUS PURPURASCENS, Pers.—In same situation as No. 58.
60. KNEIFFIA SETIGERA, Fr.—Occasional on fallen limbs.
61. CORTICIUM POLYGNII, Fr.—On fallen limbs. Well marked. Not common.
62. CORTICIUM ALUTARIUM, B. & C.—Common at times in dry woods.
63. CORTICIUM OCHROLEUCUM, Fr.—On old fallen limbs and trees, under side, in contact with the earth. White. Velvety in texture. May be the same as *C. radians*, Berk., an unpublished species from Venezuela, and Mr. Ellis thinks it is *C. radiosum*, Fr.
64. CORTICIUM OCHROLEUCUM, Fr., var.—Distinct from No. 63 by its ochre-red color, which is well defined in every specimen. It should be called a good species.
65. CORTICIUM ARACHNOIDEUM, B.—Abundant on fallen limbs.
66. CORTICIUM LACTESCENS, B.—Very common. White, milky.
67. CORTICIUM PORTENTOSUM, B. & C.—White, thick, under side of old logs. Resembles No. 63. This is a tropical species.
68. CORTICIUM INCARNATUM, Fr.—Common.
69. CORTICIUM DENDROIDEUM, n. sp. MS.—Found on a small limb. Color brownish-yellow.
70. CORTICIUM CERVICOLOR, B. & C.—No. 20 and 150 of Ellis, who thinks it may yet prove to be a new genus. Very rare and fine.
71. CORTICIUM SULFUREUM, Fr.
72. CORTICIUM DRYINUM, B. & C.—On fallen bark. Rare.
73. CORTICIUM GIGANTEUM, Fr.—Not common. Much like *C. subgiganteum*, B. & C., *læve*, Pers.
74. CORTICIUM SUBGIGANTEUM, B.—Not uncommon.
75. CORTICIUM LÆVE, Pers.—Abundant on limbs. Hard to separate from 73 and 74.
76. CORTICIUM PETERSII, B. & C.—On rotten wood. Abundant.
77. CORTICIUM GLABRUM, B. & C.—On dead bark.
78. CORTICIUM SCUTELLARE, B. & C.—Well marked. On limbs.
- It appears that *Corticium* abounds in species. There may be others, but we are certain of the foregoing, and their elimination has fully occupied the time of Mr. E., myself and others. They will receive, however, further study, and the results will be noted.

NOTE AND CORRECTION.

It is stated on page 87 in connection with the collection made by the Rev. A. B. Langlois that Plaquemines county is not far from Baton Rouge. Mr. Langlois informs us that this is true of Plaquemines city, but that Pointe a la Hache (his P. O. address), the county seat of Plaquemines county, is 200 miles from Baton Rouge, below New Orleans. If Plaquemines is not at all mentioned in his address, no confusion would arise.

On page 54, 24,000 should read 2,400.